

## Do fungal spore morphological traits correlate with allergenicity?

Raphael R. Sanvictores<sup>1</sup>, Michelle Charina C. Gomez<sup>1</sup>, Stephen Louie R. Briones<sup>1</sup>, Wan Teng M. Chang<sup>1</sup>, Adrienne Marie A. Morales<sup>1</sup>, Nikki Heherson A. Dagamac<sup>1</sup>, Thomas Edison E. dela Cruz<sup>1, 2, \*</sup>

<sup>1</sup>Department of Biological Sciences, College of Science, University of Santo Tomas, España Blvd. 1015 Manila, Philippines. <sup>2</sup>Fungal Biodiversity, Ecogenomics and Systematics (FBeS) Group, Research Center for the Natural and Applied Sciences, University of Santo Tomas, España Blvd. 1015 Manila, Philippines.

\*Correspondence: tedelacruz@ust.edu.ph

---

### Abstract

Fungal spores, like pollen, are identified as sources of allergens, yet fungal spore morphology, unlike pollen morphology, has not been correlated with allergenicity. In this study, we listed allergenic fungi reported from published literature and gathered information about their spore morphologies including the species' lifestyle, the ability to produce mycotoxins, and the types of hypersensitivity reactions they induced. We tested the association of these spore traits with the hypersensitivity reaction through correspondence analysis with Chi-square as the measure of distance. Our research listed a total of 158 species of allergenic fungi belonging to 82 genera and 30 taxonomic orders. Most of the species (n = 122) elicited a Type I hypersensitivity reaction while 33 species had more than one hypersensitivity type (Types I-III-IV). The most common allergenic fungi belong to the genus *Alternaria* (41 species). Two fungal taxa commonly found in spoiled food, *Penicillium* (9 species) and *Aspergillus* (8 species), were also listed as allergenic. We did not find any strong correlation between allergenic reaction with the following spore traits: shape, texture, color, size, appendages, and with the type of spores, presence of mycotoxins, and the species lifestyle. However, spore length and width were positively associated with hypersensitivity reaction. Allergenic fungi with short and/or narrow spores can likely cause multiple types of hypersensitivity reactions while fungi with large and/or wide spores can induce either Type I or Type III hypersensitivity reaction. Our research study provides interesting insights into the role of fungal spore morphologies in human allergenicity.

**Keywords:** allergens, allergenic fungi, hypersensitivity reaction, spore morphology, systematic review

---

Sanvictores RR, Gomez MCC, Briones SLR, Chang WTM, Morales AMA, Dagamac NHA, dela Cruz TEE (2022) Do fungal spore morphological traits correlate with allergenicity? MycoAsia 2022/05.

Received: 05.04.2022 | Accepted: 20.05.2022 | Published 20.05.2022

Handling Editor: Dr. Belle Damodara Shenoy

Reviewers: Dr. Prasanna Honnavar, Dr. Krystle Angelique Santiago

---

### Introduction

The prevalence of allergies is increasing globally, thereby affecting millions of individuals and approximately 80 % of families (Sánchez-Borges et al. 2018). Its cause ranges from dust, pollens to fungal spores, exposure to chemicals from plants and secretions from animals and insects to drugs, and ingestion of food allergens such as nuts and grains (Schnyder and Pichler 2009, Woodfolk et al. 2015, Goodman et al. 2020). From these allergens, 40 % of the world population is sensitized to pollen, leading to the common allergic rhinitis (Depciuch et al. 2016). Pollen morphology has been correlated with allergenicity, with the smaller pollen size being significant in causing the allergenic reaction. Their small size provides a greater advantage for wind dispersal, thereby increasing the pollen concentration in the air and the possibility of entering the human airways (Diethart et al. 2007, Pablos et al. 2016). In the Philippines, Sabit et al. (2020) stated that pollen allergens affect about 20 % of the Philippine population.

Another common allergen is the fungal spores. Fungi are the most abundant microorganisms in the atmosphere as their spores are easily dispersed and could arise from events and environments associated with human activities (Ortega-Rosas et al. 2019). Interestingly, the quantity of fungal spores in the atmosphere exceeds that of pollen by a thousand-fold and it can remain prevalent in the air throughout the year with seasonal peaks (Burge 2002, Boddy 2016). Fungal spores are generally smaller than pollen, making it more possible to reach the host's alveoli, and subsequently cause severe allergic reactions (Żukiewicz-Sobczak 2013). Other hypersensitivity reactions can also be triggered through ingestion or contact with these fungal spores (Simon-Nobbe et al. 2007). Since both indoor and outdoor fungi are involved in fungi-induced allergies, more than 80 fungal genera have been so far identified as allergenic which mainly induce Type I hypersensitivity (Kurup et al. 2000), albeit others can cause clinical manifestations based on Type III or Type IV hypersensitivity (Simon-Nobbe et al. 2007). However, unlike pollen, investigations on the potential contribution of fungal spore morphology to allergenicity remain scarce. This study, therefore, aims to correlate fungal spore morphology with allergenicity. We seek to delineate common fungal spore traits that may contribute to allergic reactions and serve as indicators of hypersensitivity-related risk assessment.

## **Materials and methods**

### ***Listing of allergenic fungi***

We conducted a literature survey and systematic analysis in this study. We initially utilized the paper of Simon-Nobbe et al. (2007) as the primary reference to list the species of allergenic fungi (n = 143). We expanded the list by conducting a literature search with the keywords “allergenic fungi” and “allergenic fungal species listing” on online journal databases, e.g., JSTOR (jstor.org), the National Center for Biotechnology Information (ncbi.nlm.nih.gov), Science Direct (sciencedirect.com). From our search hits, non-related studies were excluded based on information on their titles and abstracts, and on the accessibility of the paper. We found twelve studies in addition to the primary reference of Simon-Nobbe et al. (2007) to list down the allergenic fungal species, i.e., Horner et al. (1995), Green et al. (2003), Kurup (2003), Knutsen et al. (2012), Cramer et al. (2013), Kochar et al. (2014), Oh et al. (2014), Fukutomi and Taniguchi (2015), Vijayakumar et al. (2017), Bhattacharya et al. (2018), Dey et al. (2018), and Songnuan et al. (2018). In case only the fungal genus is identified in the published study, an additional literature search on the same online databases was conducted with the genus name coupled with the keywords “allergenic species” or “allergy”. Species names of the listed allergenic fungi were validated with Index Fungorum (2022).

### ***Listing of fungal spore traits***

We initially listed the set of spore morphologies and other characters for this study. These included (1) species lifestyle, (2) spore shape, (3) spore texture, (4) spore color, (5) spore size, (6) spore length, (7) spore width, (8) the type of spores, (9) the presence of spore appendages, and (10) the ability to produce mycotoxins. To determine the morphological characters for each species, we initially searched the traits from reference taxonomic monographs, i.e., the Pictorial Atlas of Soil and Seed Fungi: Morphologies of Cultured Fungi and Key to Species (Watanabe 2010) and Larone's Medically Important Fungi: A Guide to Identification (Walsh et al. 2018). Additionally, we also searched online taxonomic databases, e.g., MycoBank (mycobank.org), Mycology Online at the University of Adelaide (adelaide.edu.au/mycology), and Doctor Fungus of Mycoses Study Group Education & Research Consortium (MSG ERC) (drfungus.org). In these databases, characters were searched by typing the species name. For other missing characters, we also conducted an extensive literature search by typing the species name coupled with the following keywords: ‘spore morphology’, ‘species description’, ‘mycotoxins’, ‘lifestyle’, and ‘nutrition mode’.

Following listing of spore morphologies and other traits, these were categorized as follows: (1) *lifestyle*: biotrophic, endophytic, hemibiotrophic, necrotrophic, saprotrophic, and varying lifestyle for

species that exhibits two or more lifestyles, following the definition of De Silva et al. (2016); (2) *spore shape*: elliptical [ellipsoidal or cylindrical appearance], fusiform [spindle-like shape], globose [spherical], ovate [oval outline], quadrilateral [four-sided figure], and variable [multiple shapes]; (3) *spore texture*: smooth [partially or completely smooth], rough [has a specific ornamentation], and smooth to rough; (4) *spore color*: light, dark, and light to dark; (5) *spore size*: I [long and wide], II [long and narrow], III [short and wide], and IV [short and narrow]; (6) *spore length*: long [ $>20\ \mu\text{m}$ ] and short [ $\leq 20\ \mu\text{m}$ ]; (7) *spore width*: wide [ $>10\ \mu\text{m}$ ] and narrow [ $\leq 10\ \mu\text{m}$ ]; (8) *type of spore*: asexual spore and sexual spore; (9) *presence of spore appendages*: absent and present; and (10) *ability to produce mycotoxins*: absent and present.

### **Identifying the types of hypersensitivity reaction (HR)**

We also listed the type/s of hypersensitivity reactions for each of the listed allergenic fungi. The types follow Gell and Coombs's classification of hypersensitivity reaction for fungal allergies as described in the papers of Simon-Nobbe et al. (2007) and Sheldon et al. (2014). This is also supplemented by an extensive literature search to identify the type of hypersensitivity reaction elicited by each listed species. Species without any reported type of hypersensitivity reaction were excluded from the analysis and designated as potentially allergenic fungi.

### **Data analysis**

From our listing of species, spore traits, and hypersensitivity reactions, we initially generated a table with the frequencies per category using the PivotTable tool of Microsoft Excel Version 2202. We did correspondence analysis (CA) in row principal normalization with Chi-square ( $\chi^2$ ) as the distance measure with IBM SPSS Statistics version 20. The spore traits were designated as the row variables while the types of hypersensitivity reaction were assigned as the column variables. For each character, statistical values from the  $\chi^2$  test of independence at a 5 % significance level, the Cramér's *V* coefficient, and a biplot were generated. To interpret the association patterns, spore traits within the same quadrant of a particular hypersensitivity reaction are deemed strongly associated. For spore traits with only two categories, a multiple correspondence analysis (MCA) with symmetrical normalization with  $\chi^2$  as the distance measure was conducted.

### **Results**

**Allergenic and potentially allergenic fungi.** A total of 33 taxonomic orders, 101 genera, and 192 species of allergy-associated fungi were listed based on our literature analysis. Of these, 30 taxonomic orders (excluding Dacrymycetales, Geastrales, and Peronosporales) and 82 genera were confirmed as allergenic as indicated by their reported hypersensitivity reactions (Table 1). The excluded taxonomic orders had these three species, *Dacrymyces deliquescens* (Dacrymycetales), *Geastrum saccatum* (Geastrales), and *Plasmopara viticola* (Peronosporales), identified as potentially allergenic species but without any reported hypersensitivity reactions. We also identified 24 species as potentially allergenic as these were listed as allergenic fungi in published literature but with no reported hypersensitivity reactions (Table 3). Of these 24 taxa, 15 were identified only at the genus level, and hence, their species identity could not be ascertained. In addition, 10 species were reported as allergenic with Type I hypersensitivity reaction, but identification was limited only to the genus level (Table 3). Interestingly, two genera have both reported allergenic (AF) and potentially allergenic (PAF) fungal species – *Aspergillus* (8 AF, 4 PAF) and *Penicillium* (9 AF, 1 PAF). Of the 192 species reported from published literature, 158 were identified as allergenic, with the majority exhibiting Type I hypersensitivity reaction (Table 1). The genus *Alternaria* had the highest number of allergenic species, a total of 41 species (Table 2). This is followed by *Aspergillus* and *Penicillium*, with 8 and 9 identified species, respectively. The top ten genera with the greatest number of recorded allergenic fungi are shown in Table 2.

**Table 1.** The number of taxa per type of hypersensitivity reaction

|         | Type I | Type III | Multiple HR Types <sup>a</sup> | Total |
|---------|--------|----------|--------------------------------|-------|
| Order   | 15     | 0        | 15                             | 30    |
| Genus   | 56     | 2        | 24                             | 82    |
| Species | 122    | 3        | 33                             | 158   |

<sup>a</sup>Multiple hypersensitivity reaction (HR) Types: Type I + III, Type I + IV, Type III + IV, Type I + III + IV

**Table 2.** The number of species recorded for the top 10 allergenic fungi

| Genera              | Number of Allergenic Fungal Species |
|---------------------|-------------------------------------|
| <i>Alternaria</i>   | 41                                  |
| <i>Penicillium</i>  | 9                                   |
| <i>Aspergillus</i>  | 8                                   |
| <i>Curvularia</i>   | 5                                   |
| <i>Fusarium</i>     | 5                                   |
| <i>Calvatia</i>     | 3                                   |
| <i>Cladosporium</i> | 3                                   |
| <i>Mucor</i>        | 3                                   |
| <i>Stemphylium</i>  | 3                                   |
| <i>Trichophyton</i> | 3                                   |

**Table 3.** List of fungal species with uncertain allergenicity and identities

**Potentially allergenic fungi:** species reported as allergenic in published literature but with no reported type of hypersensitivity reaction

|                                |                                  |                         |
|--------------------------------|----------------------------------|-------------------------|
| <i>Aspergillus flavipes</i>    | <i>Laetiporus</i> sp.            | <i>Torula</i> sp.       |
| <i>A. giganteus</i>            | <i>Microsporium</i> sp.          | <i>Verticillium</i> sp. |
| <i>A. sydowii</i>              | <i>Monilinia</i> sp.             | <i>Xylaria</i> sp.      |
| <i>A. ustus</i>                | <i>Paecilomyces</i> sp.          | <i>Xylobolus</i> sp.    |
| <i>Botryotrichum</i> sp.       | <i>Penicillium solitum</i>       |                         |
| <i>Choanephora</i> sp.         | <i>Pithomyces</i> sp.            |                         |
| <i>Circinella</i> sp.          | <i>Plasmopara viticola</i>       |                         |
| <i>Cyathus</i> sp.             | <i>Pyrenochaeta</i> sp.          |                         |
| <i>Dacrymyces deliquescens</i> | <i>Spegazzinia</i> sp.           |                         |
| <i>Geastrum saccatum</i>       | <i>Syncephalastrum racemosum</i> |                         |

**Allergenic fungi:** species reported as allergens with the reported types of hypersensitivity reactions in published literature but with no species identity

|                           |                           |                           |
|---------------------------|---------------------------|---------------------------|
| <i>Chaetomium</i> sp.     |                           | Simon-Nobbe et al. (2007) |
| <i>Claviceps</i> sp.      |                           | Simon-Nobbe et al. (2007) |
| <i>Cylindrocarpon</i> sp. |                           | Simon-Nobbe et al. (2007) |
| <i>Erysiphe</i> sp.       |                           | Kurup (2003)              |
| <i>Spondycladium</i> sp.  | Type I                    | Burrows et al. (1989)     |
| <i>Sporobolomyces</i> sp. | Hypersensitivity Reaction | Simon-Nobbe et al. (2007) |
| <i>Tilletia</i> sp.       |                           | Simon-Nobbe et al. (2007) |
| <i>Tilletiopsis</i> sp.   |                           | Simon-Nobbe et al. (2007) |
| <i>Trichothecium</i> sp.  |                           | Simon-Nobbe et al. (2007) |
| <i>Urocystis</i> sp.      |                           | Kurup (2003)              |

**Lifestyle.** Most of the allergenic fungi had varying life strategies from saprotrophic to biotrophic and endophytic and elicit a Type I hypersensitivity reaction (Table 4). However, our statistical analysis showed no relationship between the different lifestyles and the types of hypersensitivity reactions (Table 5, Figure 1), and the level of association was found to be negligible (Table 6).


**Spore shape, texture, and color.** The most prominent shape among the listed allergenic species was elliptical, followed by globose, ovate, fusiform, and quadrilateral (Table 4). Many species also exhibited variable shapes. Spore texture is generally smooth (92 species), albeit half of this number equivalent to 46 species had a rough texture. For spore color, the number of species is almost equally distributed between light- and dark-colored spores and a combination of both. A greater number of our listed species had Type I hypersensitivity reaction in relation to these spore traits, albeit we did not observe any strong relationship (Table 5, Figure 1), The Cramér's *V* coefficient also suggests a negligible to a weak association between the hypersensitivity reactions and these spore traits (Table 6).

**Spore size, length, and width.** Spore size, i.e., length plus width (Table 4), was mostly short and narrow (category IV, 93 species) followed by long and wide (Category I, 48 species). The observed  $\chi^2$  value, the significance probability, and the Cramér's *V* coefficient all showed no to a weak association between the type of hypersensitivity reaction elicited by the examined species and the overall spore size (Tables 5–6, Figure 1).

**Table 4.** Frequency of allergenic fungi per spore traits and types of hypersensitivity reaction

| Characters                    | Categories                         | Type I | Type III | Multiple HR Types <sup>a</sup> | Total |
|-------------------------------|------------------------------------|--------|----------|--------------------------------|-------|
| Lifestyle                     | Biotrophic                         | 4      | 0        | 0                              | 4     |
|                               | Endophytic                         | 24     | 1        | 6                              | 31    |
|                               | Saprotrophic                       | 28     | 1        | 10                             | 39    |
|                               | Varying Lifestyle                  | 70     | 1        | 15                             | 86    |
| Spore Shape                   | Elliptical Shape                   | 42     | 1        | 8                              | 51    |
|                               | Fusiform Shape                     | 5      | 0        | 1                              | 6     |
|                               | Globose Shape                      | 21     | 1        | 5                              | 27    |
|                               | Ovate Shape                        | 5      | 1        | 1                              | 7     |
|                               | Quadrilateral Shape                | 1      | 0        | 1                              | 2     |
|                               | Variable Shape                     | 59     | 0        | 16                             | 75    |
| Spore Texture                 | Smooth                             | 69     | 2        | 21                             | 92    |
|                               | Rough                              | 37     | 1        | 8                              | 46    |
|                               | Smooth to Rough                    | 21     | 0        | 2                              | 23    |
| Spore Color                   | Dark Color                         | 47     | 1        | 9                              | 57    |
|                               | Light Color                        | 43     | 1        | 14                             | 58    |
|                               | Light to Dark Color                | 40     | 1        | 7                              | 48    |
| Spore Size<br>(Length, Width) | I (Long, Wide)                     | 44     | 0        | 4                              | 48    |
|                               | II (Long, Narrow)                  | 12     | 0        | 2                              | 14    |
|                               | III (Short, Wide)                  | 10     | 0        | 1                              | 11    |
|                               | IV (Short, Narrow)                 | 65     | 3        | 25                             | 93    |
| Spore Length                  | Long (>20 $\mu\text{m}$ )          | 56     | 0        | 6                              | 62    |
|                               | Short ( $\leq 20$ $\mu\text{m}$ )  | 75     | 3        | 26                             | 104   |
| Spore Width                   | Wide (>10 $\mu\text{m}$ )          | 54     | 0        | 5                              | 59    |
|                               | Narrow ( $\leq 10$ $\mu\text{m}$ ) | 77     | 3        | 27                             | 107   |
| Type of Spore                 | Asexual Spore                      | 97     | 2        | 24                             | 123   |
|                               | Sexual Spore                       | 36     | 1        | 8                              | 45    |
| Presence of Appendages        | Appendages Absent                  | 127    | 3        | 31                             | 163   |
|                               | Appendages Present                 | 4      | 0        | 1                              | 5     |
| Presence of Mycotoxins        | Mycotoxins Absent                  | 72     | 1        | 18                             | 91    |
|                               | Mycotoxins Present                 | 61     | 2        | 14                             | 77    |

<sup>a</sup>Multiple hypersensitivity reaction (HR) Types: Type I + III, Type I + IV, Type III + IV, Type I + III + IV



**Figure 1.** Association patterns between the types of hypersensitivity reaction and (A) lifestyle, (B) spore shape, (C) spore texture, (D) spore color, and (E) spore size of allergenic fungi

When spore size was categorized separately as length or width, the majority of the allergenic fungi exhibited a short length and a narrow width or diameter regardless of their hypersensitivity reaction (Table 4). Interestingly, while we did not observe any association between spore size and hypersensitivity reaction, there is a moderate association between spore length and the types of hypersensitivity reaction as elicited by the allergenic fungi (Tables 5–6). The same moderate association was also observed between spore width and HR types. Interestingly, the MCA plot showed that a short spore length is most likely associated with Type III and multiple types of hypersensitivity reactions while a long spore length is probably associated with Type I (Figure 2). Similarly, the MCA plot also showed that allergenic spores with a narrow width are likely associated with Type III and multiple types of hypersensitivity reactions while allergenic spores with a wide width may have an association with inducing a Type I hypersensitivity reaction.

**Other Characters.** Most of the listed allergenic fungi have asexual spores. Their spores also had no appendages. The fungi also had no reported mycotoxins. And just like the other spore traits, these species' characteristics showed negligible association with the types of hypersensitivity reactions (Tables 4–6, Figure 2).

**Table 5.**  $\chi^2$  test of independence between the spore traits and types of hypersensitivity reaction


| Characters             | $\chi^2$<br>(Observed value) | $\chi^2$<br>(Critical value) | P<br>( $\alpha = .05$ ) |
|------------------------|------------------------------|------------------------------|-------------------------|
| Lifestyle              | 2.992                        | 12.592                       | 0.810                   |
| Spore Shape            | 10.147                       | 18.307                       | 0.428                   |
| Spore Texture          | 3.164                        | 9.488                        | 0.531                   |
| Spore Color            | 2.009                        | 9.488                        | 0.734                   |
| Spore Size             | 11.131                       | 12.592                       | 0.084                   |
| Spore Length           | 8.151                        | 5.991                        | 0.017                   |
| Spore Width            | 9.039                        | 5.991                        | 0.011                   |
| Type of Spores         | 0.123                        | 5.991                        | 0.940                   |
| Presence of Appendages | 0.095                        | 5.991                        | 0.954                   |
| Presence of Mycotoxins | 0.580                        | 5.991                        | 0.748                   |

**Table 6.** Cramér's V coefficient for each of the spore traits of the allergenic fungi

| Characters             | Cramér's V | Interpretation of Association <sup>a</sup> |
|------------------------|------------|--|
| Lifestyle              | 0.096      | Negligible                                 |
| Spore Shape            | 0.174      | Weak                                       |
| Spore Texture          | 0.099      | Negligible                                 |
| Spore Color            | 0.079      | Negligible                                 |
| Spore Size             | 0.183      | Weak                                       |
| Spore Length           | 0.222      | Moderate                                   |
| Spore Width            | 0.233      | Moderate                                   |
| Type of Spores         | 0.027      | Negligible                                 |
| Presence of Appendages | 0.024      | Negligible                                 |
| Presence of Mycotoxins | 0.059      | Negligible                                 |

<sup>a</sup>Association Values: 0 – 0.100 (negligible), 0.101 – 0.200 (weak), 0.201 – 0.400 (moderate), 0.401 – 0.600 (relatively strong), 0.601 – 0.800 (strong), 0.801 – 1.000 (very strong)

**Spore traits of potentially allergenic fungi.** These fungal species were reported in the published literature as allergenic species. However, there are no published reports on the types of hypersensitivity reactions elicited by these species, and hence, we categorized these fungi here as potentially allergenic. Given the association we observed between spore length and width and HR types, we predicted the probable types of hypersensitivity reactions that these potentially allergenic fungi will probably exhibit (Table 7). But it is important to note that laboratory tests must be conducted to verify these predictions.



**Figure 2.** Association patterns between the types of hypersensitivity reaction and the (A) spore length and width, and (B) other traits of allergenic fungi

**Table 7.** Predicted types of hypersensitivity reactions for the listed potentially allergenic fungi based on the association patterns with spore length and width

| Genus                  | Species                  | Length | Width  | Predicted HR Type(s)                    |
|------------------------|--------------------------|--------|--------|---|
| <i>Aspergillus</i>     | <i>A. flavipes</i>       | Short  | Narrow | Type III or Multiple HR types           |
|                        | <i>A. giganteus</i>      | Short  | Narrow | Type III or Multiple HR types           |
|                        | <i>A. sydowii</i>        | Short  | Narrow | Type III or Multiple HR types           |
|                        | <i>A. ustus</i>          | Short  | Narrow | Type III or Multiple HR types           |
| <i>Botryotrichum</i>   | <i>Botryotrichum</i> sp. | Short  | Wide   | Type I or Type III or Multiple HR types |
| <i>Choanephora</i>     | <i>Choanephora</i> sp.   | Short  | Narrow | Type III or Multiple HR types           |
| <i>Circinella</i>      | <i>Circinella</i> sp.    | Short  | Narrow | Type III or Multiple HR types           |
| <i>Cyathus</i>         | <i>Cyathus</i> sp.       | Short  | Narrow | Type III or Multiple HR types           |
| <i>Dacrymyces</i>      | <i>D. deliquescens</i>   | Short  | Narrow | Type III or Multiple HR types           |
| <i>Geastrum</i>        | <i>G. saccatum</i>       | Short  | Narrow | Type III or Multiple HR types           |
| <i>Laetiporus</i>      | <i>Laetiporus</i> sp.    | Short  | Narrow | Type III or Multiple HR types           |
| <i>Microsporium</i>    | <i>Microsporium</i> sp.  | Long   | Wide   | Type I                                  |
| <i>Monilinia</i>       | <i>Monilinia</i> sp.     | Short  | Wide   | Type I or Type III or Multiple HR types |
| <i>Paecilomyces</i>    | <i>Paecilomyces</i> sp.  | Short  | Narrow | Type III or Multiple HR types           |
| <i>Penicillium</i>     | <i>P. solitum</i>        | Short  | Narrow | Type III or Multiple HR types           |
| <i>Pithomyces</i>      | <i>Pithomyces</i> sp.    | Long   | Wide   | Type I                                  |
| <i>Plasmopara</i>      | <i>P. viticola</i>       | Short  | Narrow | Type III or Multiple HR types           |
| <i>Pyrenochaeta</i>    | <i>Pyrenochaeta</i> sp.  | Short  | Narrow | Type III or Multiple HR types           |
| <i>Spegazzinia</i>     | <i>Spegazzinia</i> sp.   | Short  | Wide   | Type I or Type III or Multiple HR types |
| <i>Syncephalastrum</i> | <i>S. racemosum</i>      | Short  | Narrow | Type III or Multiple HR types           |
| <i>Torula</i>          | <i>Torula</i> sp.        | Short  | Narrow | Type III or Multiple HR types           |
| <i>Verticillium</i>    | <i>Verticillium</i> sp.  | Short  | Narrow | Type III or Multiple HR types           |
| <i>Xylaria</i>         | <i>Xylaria</i> sp.       | Short  | Narrow | Type III or Multiple HR types           |
| <i>Xylobolus</i>       | <i>Xylobolus</i> sp.     | Short  | Narrow | Type III or Multiple HR types           |



## Discussion

Pollen morphology has been associated with pollen allergenicity. The typical aerodynamic size of pollen, which ranges from 15–40  $\mu\text{m}$ , allows for its deposition on the nasopharynx (D'Amato et al. 2007). There are also pieces of evidence that pollen could enter the lower respiratory tract, and hence, is implicated in asthma (Busse et al. 1972, D'Amato et al. 2007). Pollen that is associated with allergenicity lacks a distinguishable endexine layer in its pollen wall, which results in the leakage of allergenic proteins (Diethart et al. 2007). Like pollen, fungal spores have also been implicated in triggering hypersensitivity reactions among susceptible individuals. While fungal spores generally function in the species dispersal in different habitats and in species survival under unfavorable conditions (Dijksterhuis 2019), fungal spores when inhaled can cause hypersensitivity reactions as it contains immunoglobulin-specific antigens (Horner et al. 1995). In our systematic review, 158 fungal species were identified as capable of eliciting an allergic reaction. An additional 24 species were also listed as allergenic fungi but do not have any reported hypersensitivity reaction. This led us to look at spore traits and if any of its morphological properties are associated with a specific hypersensitivity reaction to predict the possible allergenicity of these potentially allergenic fungi.

Certain morphological traits may be factors to the fungal allergenicity. For example, spore shape can influence spore release, wherein an elliptical shape maximizes flight speed (Roper et al. 2008, Halbwachs and Bässler 2021), which in turn can be correlated with greater spore dispersal and persistence in the atmosphere. A globose spore shape can reduce the surface area to volume ratio, thereby facilitating better survival for the organism (Bergman and Casadevall 2010, Wang and Lin 2012, Calhim et al. 2018). Smooth spores or spores with simple ornamentation also allow for better wind dispersal (Diethart et al. 2007, Dawson et al. 2018) while the presence of spiny features on the spore surface can reduce contact, resulting in lower adherence (Whitehead et al. 2021). For spore color, pigments produced by fungi protect species from environmental stressors such as UV (Horner et al. 1995). Spore appendages can also function for attachment to a preferred host as observed with the spore dispersal mechanism of marine and freshwater fungi (Crous et al. 2012). However, unlike pollen morphology, these spore traits were not systematically tested for their association with hypersensitivity reactions.

In addition to spore traits, we also tested the correlation of lifestyles, type of spores, and mycotoxin production with allergenicity. Fungi are known to utilize various life strategies (i.e., biotrophic, endophytic, hemibiotrophic, necrotrophic, saprotrophic, or a combination of these) to facilitate adaptation in a changing environment (Girard et al. 2013, Muszewska et al. 2017). Asexual spores are produced in extensive numbers, ensuring that the organism can be quickly dispersed. Additionally, asexual spores serve as a resistance mechanism against harsh environmental conditions (Peberdy 1980). In contrast, the production of sexual spores increases genetic diversity, and when conditions are constantly fluctuating may select for the better-adapted strains (Sun and Heitman 2011). Mycotoxins also play a role in the virulence and pathogenicity of fungi, with the adverse effects being dependent on the level of exposure (Bush et al. 2006, Simon-Nobbe et al. 2007). Mycotoxins, e.g., gliotoxin and patulin, were considered aggravators of allergic asthma in murine models (Schütze et al. 2010, Kraft et al. 2021). Despite the potential of these spore traits (type, shape, color, texture, presence of appendages) and other species features (lifestyle, mycotoxin production) for dispersal, persistence, and survival of fungi, we did not find any strong association between these morphological features with any type of hypersensitivity reaction (Tables 5–6). However, we found a moderate association between spore size, particularly spore length and spore width, and types of hypersensitivity reaction (Tables 5–6, Figure 2).

A link between the size of the aeroallergens and the anatomical site to which these aeroallergens can be deposited is already established. For example, large spores are correlated with the upper respiratory

airways, thereby causing allergic diseases in the nasopharyngeal area and the paranasal sinuses (D'Ovidio et al. 2021). In contrast, small spores were found to penetrate the lower respiratory tract and trigger allergic diseases of the bronchi and alveoli (Yamamoto et al. 2012, Dey et al. 2018). This link can be attributed to the spore deposition mechanisms in the respiratory tract, i.e., impaction and sedimentation (Deacon 2006).

The main mechanism, impaction, by which large spores are deposited, is observed during airflow fluctuation. This fluctuation causes the spores to diverge from their initial trajectories and impact the walls of the respiratory airways (Darquenne 2020). This only occurs in the upper respiratory airways since the structure of the nasal cavity increases air turbulence (Marieb and Hoehn 2018). Additionally, the size contributes to the momentum of large spores to impact and become trapped in the nasal mucosa (Deacon 2006). This elucidated why most large spores are associated with allergic sinusitis (Type I and/or Type III) and allergic rhinitis (Type I) (Simon-Nobbe et al. 2007, Ryan and Clark 2015). On the other hand, small size is insufficient for the spores to impact but enable the spores to reach the lower respiratory tract when the airflow is laminar (Deacon 2006). The primary mechanism for small spores is sedimentation, wherein the buoyancy of particles is reduced by gravitational forces (Darquenne 2020). In other words, the suspended small spores are acted upon by gravity, allowing them to finally settle on the surface or walls of the lower respiratory airways. Such phenomenon explains why small spores can induce multiple types of hypersensitivity reactions as manifested by allergic diseases such as hypersensitivity pneumonitis (Type III and Type IV), allergic bronchopulmonary mycosis (Type I, Type III, and Type IV), and the exacerbation of allergic asthma (Type I) (Mohr 2004, Simon-Nobbe et al. 2007, Agarwal and Gupta 2011, Shah and Panjabi 2014, Riario-Sforza and Marinou 2017, Pfavayi et al. 2020). Interestingly, some of the listed species with spores that are not entirely small in our review were implicated in allergic diseases affecting the lower respiratory tract. Sensitization to *Alternaria* and *Curvularia* were associated with severe allergic asthma, possibly due to the continuous release of allergens from the upper airways (Denning et al. 2014).

To confirm the types of hypersensitivity reactions induced by allergenic fungi, *in vitro* assays can be performed to assess the sensitizing potential of a spore. These assays include the radioallergosorbent test (RAST) which is specific to a type of immunoglobulin, the basophil activation test (BAT), the nephelometric detection of immunoglobulins or immune complexes, the radial immunodiffusion (RID), the lymphocyte transformation test (LTT), and the electro immunoassay (Virella et al. 1979, Whicher et al. 1984, Naisbitt et al. 2014, Foong et al. 2021). The enzyme-linked immunosorbent assay (ELISA) and Western blot analysis have also been used (Simon-Nobbe et al. 2007). To supplement the findings from these methods, *in vivo* allergy testing, specifically skin prick testing, is taken together with a history of the sensitized patient (Foong et al. 2021). We believe that spore size may also help with identifying potential allergenic fungi as an important preliminary step. In this study, we identified the possible type of hypersensitivity reactions that can be induced by potential allergenic fungi based on their spore length and spore width (Table 7). However, to confirm this, any of the *in vitro* assays must be conducted to validate our prediction.

In conclusion, following our systematic review, we found out that allergenic fungi with small spores are most likely able to trigger multiple types or a combination of hypersensitivity reactions. Fungal species with large spores, which can be associated with Type I hypersensitivity reaction, may be considered less potent to cause allergic diseases due to the limitations posed by the narrowing respiratory tract and deposition mechanisms. Spore morphological traits can offer insights into the allergenicity of fungi and can be utilized as a preliminary basis for hypersensitivity-related risk assessment of fungal species. Future studies may also focus on correlating spore morphological traits

of other spore-bearing organisms such as myxomycetes (slime molds) to types of hypersensitivity reactions and/or specific allergic diseases.

### Acknowledgments

The authors acknowledge all the authors whose published works served as the basis for this study.

### Statement on conflict of interest

The authors declare no conflict of interest.

### Authors contribution

TEEDC conceptualized the research study. RRS, MCCG, SLRB, WTMC, and AMAM conducted the literature survey and data collection. RRS, MCCG, SLRB, WTMC, AMAM, NHAD, and TEEDC analyze the data and wrote the manuscript. The authors contributed equally to this work.

### References

- Agarwal R, Gupta D (2011) Severe asthma and fungi: current evidence. *Medical Mycology* 49:S150-S157. <https://doi.org/10.3109/13693786.2010.504752>
- Bergman A, Casadevall A (2010) Mammalian endothermy optimally restricts fungi and metabolic costs. *MBio* 1(5): e00212-10. <https://doi.org/10.1128/mbio.00212-10>
- Bhattacharya K, Sircar G, Dasgupta A, Gupta Bhattacharya S (2018) Spectrum of allergens and allergen biology in India. *International Archives of Allergy and Immunology* 177:219-237. <https://doi.org/10.1159/000490805>
- Boddy L (2016) Interactions with humans and other animals. In: Watkinson SC, Boddy L, Money NP (eds.) *The Fungi*. Academic Press, 293-336. <https://doi.org/10.1016/B978-0-12-382034-1.00009-8>
- Burge H (2002) An update on pollen and fungal spore aerobiology. *Journal of Allergy and Clinical Immunology* 110:544-552. <https://doi.org/10.1067/mai.2002.128674>
- Burrows B, Martinez FD, Halonen M., Barbee RA, Cline MG (1989) Association of asthma with serum IgE levels and skin-test reactivity to allergens. *New England Journal of Medicine* 320:271-277. <https://doi.org/10.1056/nejm198902023200502>
- Bush R, Portnoy J, Saxon A, Terr A, Wood R (2006) The medical effects of mold exposure. *Journal of Allergy and Clinical Immunology* 117:326-333. <https://doi.org/10.1016/j.jaci.2005.12.001>
- Busse WW, Reed CE, Hoehne, JH (1972) Where is the allergic reaction in ragweed asthma? *Journal of Allergy and Clinical Immunology* 50:289-293. [https://doi.org/10.1016/0091-6749\(72\)90027-9](https://doi.org/10.1016/0091-6749(72)90027-9)
- Calhim S, Halme P, Petersen J, Læssøe T, Bässler C, Heilmann-Clausen J (2018) Fungal spore diversity reflects substrate-specific deposition challenges. *Scientific Reports* 8(1):5356. <https://doi.org/10.1038/s41598-018-23292-8>
- Crameri R, Garbani M, Rhyner C, Huitema C (2013) Fungi: the neglected allergenic sources. *Allergy* 69:176-185. <https://doi.org/10.1111/all.12325>
- Crous PW, Verkley GJM, Christensen M, Castañeda-Ruiz RF, Groenewald JZ (2012) How important are conidial appendages? *Persoonia* 28:126-137. <https://doi.org/10.3767/003158512x652624>
- D'Amato G, Cecchi L, Bonini S, Nunes C, Annesi-Maesano I, Behrendt H, Liccardi G, Popov T, Van Cauwenberge P (2007) Allergenic pollen and pollen allergy in Europe. *Allergy* 62:976-990. <https://doi.org/10.1111/j.1398-9995.2007.01393.x>
- Darquenne C (2020) Deposition mechanisms. *Journal of Aerosol Medicine and Pulmonary Drug Delivery* 33:181-185. <https://doi.org/10.1089/jamp.2020.29029.cd>
- Dawson S, Boddy L, Halbwachs H, Bässler C, Andrew C, Crowther T, Heilmann-Clausen J, Nordén J, Ovaskainen O, Jönsson M (2018) Handbook for the measurement of macrofungal functional traits: A start with basidiomycete wood fungi. *Functional Ecology* 33:372-387. <https://doi.org/10.1111/1365-2435.13239>

- Deacon J (2006) Fungal Biology. Malden, MA: Blackwell Publishing.
- Denning D, Pashley C, Hartl D, Wardlaw A, Godet C, Del Giacco S, Delhaes L, Sergejeva S (2014) Fungal allergy in asthma—state of the art and research needs. *Clinical and Translational Allergy* 4(1):14. <https://doi.org/10.1186/2045-7022-4-14>
- Depciuch J, Kasprzyk I, Roga E, Parlinska-Wojtan M (2016) Analysis of morphological and molecular composition changes in allergenic *Artemisia vulgaris* L. pollen under traffic pollution using SEM and FTIR spectroscopy. *Environmental Science and Pollution Research* 23:23203-23214. <https://doi.org/10.1007/s11356-016-7554-8>
- De Silva N, Lumyong S, Hyde K, Bulgakov T, Phillips A, Yan J (2016) Mycosphere Essays 9: Defining biotrophs and hemibiotrophs. *Mycosphere* 7:545-559. <https://doi.org/10.5943/mycosphere/7/5/2>
- Dey D, Ghosal K, Bhattacharya S (2018) Aerial fungal spectrum of Kolkata, India, along with their allergenic impact on the public health: a quantitative and qualitative evaluation. *Aerobiologia* 35:15-25. <https://doi.org/10.1007/s10453-018-9534-6>
- Diethart B, Sam S, Weber M (2007) Walls of allergenic pollen: Special reference to the endexine. *Grana* 46:164-175. <https://doi.org/10.1080/00173130701472181>
- Dijksterhuis J (2019) Fungal spores: Highly variable and stress-resistant vehicles for distribution and spoilage. *Food Microbiology* 81:2-11. <https://doi.org/10.1016/j.fm.2018.11.006>
- D'Ovidio MC, Di Renzi S, Capone P, Pelliccioni A (2021) Pollen and fungal spores evaluation in relation to occupants and microclimate in indoor workplaces. *Sustainability* 13:3154. <https://doi.org/10.3390/su13063154>
- Foong R, Dantzer J, Wood R, Santos A (2021) Improving diagnostic accuracy in food allergy. *The Journal of Allergy and Clinical Immunology: In Practice* 9:71-80. <https://doi.org/10.1016/j.jaip.2020.09.037>
- Fukutomi Y, Taniguchi M (2015) Sensitization to fungal allergens: Resolved and unresolved issues. *Allergology International* 64:321-331. <https://doi.org/10.1016/j.alit.2015.05.007>
- Girard V, Dieryckx C, Job C, Job D (2013) Secretomes: The fungal strike force. *Proteomics* 13:597-608. <https://doi.org/10.1002/pmic.201200282>
- Goodman RE, Chapman MD, Slater JE (2020) The allergen: sources, extracts, and molecules for diagnosis of allergic disease. *The Journal of Allergy and Clinical Immunology: In Practice* 8:2506-2514. <https://doi.org/10.1016/j.jaip.2020.06.043>
- Green B, Zinovia Mitakakis T, Tovey E (2003) Allergen detection from 11 fungal species before and after germination. *Journal of Allergy and Clinical Immunology* 111:285-289. <https://doi.org/10.1067/mai.2003.57>
- Halbwachs H, Bässler C (2021) Functional traits of stipitate basidiomycetes. In: Zaragoza Ó, Casadevall A (eds.) *Encyclopedia of Mycology*, 361-337. <https://doi.org/10.1016/b978-0-12-819990-9.00047-0>
- Hamilos D (2010) Allergic fungal rhinitis and rhinosinusitis. *Proceedings of the American Thoracic Society* 7:245-252. <https://doi.org/10.1513/pats.200909-098al>
- Horner W, Helbling A, Salvaggio J, Lehrer S (1995) Fungal allergens. *Clinical Microbiology Reviews* 8:161-179. <https://doi.org/10.1128/cmr.8.2.161>
- Kita T, Nishi K, Fujimura M, Abo M, Ohka T, Yasui M, Ogawa H, Minato H, Kurumaya H, Nakao S (2003) A case of hypersensitivity pneumonitis caused by *Humicola fuscoatra*. *Respirology* 8:95-98. <https://doi.org/10.1046/j.1440-1843.2003.00434.x>
- Knutsen A, Bush R, Demain J, Denning D, Dixit A, Fairs A, Greenberger P, Kariuki B, Kita H, Kurup V, Moss R, Niven R, Pashley C, Slavin R, Vijay H, Wardlaw A (2012) Fungi and allergic lower respiratory tract diseases. *Journal of Allergy and Clinical Immunology* 129:280-291. <https://doi.org/10.1016/j.jaci.2011.12.970>
- Kochar S, Ahlawat M, Dahiya P, Chaudhary D (2014) Assessment of allergenicity to fungal allergens of Rohtak City, Haryana, India. *Allergy and Rhinology* 5:e56-e65.

<https://doi.org/10.2500/ar.2014.5.0088>

- Kraft S, Buchenauer L, Polte T (2021) Mold, mycotoxins, and a dysregulated immune system: A combination of concern? *International Journal of Molecular Sciences* 22:12269. <https://doi.org/10.3390/ijms222212269>
- Kurup V (2003) Fungal allergens. *Current Allergy and Asthma Reports* 3:416-423. <https://doi.org/10.1007/s11882-003-0078-6>
- Kurup V, Shen H, Banerjee B (2000) Respiratory fungal allergy. *Microbes and Infection* 2:1101-1110. [https://doi.org/10.1016/s1286-4579\(00\)01264-8](https://doi.org/10.1016/s1286-4579(00)01264-8)
- Marieb E, Hoehn K (2018) *Human Anatomy and Physiology*. London: Pearson Education, Ltd.
- Mohr L (2004) Hypersensitivity pneumonitis. *Current Opinion in Pulmonary Medicine* 10:401-411. <https://doi.org/10.1097/01.mcp.0000135675.95674.29>
- Muszewska A, Stepniewska-Dziubinska MM, Steczkiewicz K, Pawlowska J, Dziedzic A, Ginalski K (2017) Fungal lifestyle reflected in serine protease repertoire. *Scientific Reports* 7:1-12. <https://doi.org/10.1038/s41598-017-09644-w>
- Naisbitt D, Natrass R, Ogese M (2014) *In vitro* diagnosis of delayed-type drug hypersensitivity. *Immunology and Allergy Clinics of North America* 34:691-705. <https://doi.org/10.1016/j.iac.2014.04.009>
- Oh S, Fong J, Park M, Chang L, Lim Y (2014) Identifying airborne fungi in Seoul, Korea using metagenomics. *Journal of Microbiology* 52:465-472. <https://doi.org/10.1007/s12275-014-3550-1>
- Ortega-Rosas C, Calderón-Ezquerro M, Gutiérrez-Ruacho O (2019) Fungal spores and pollen are correlated with meteorological variables: effects in human health at Hermosillo, Sonora, Mexico. *International Journal of Environmental Health Research* 30:677-695. <https://doi.org/10.1080/09603123.2019.1625031>
- Pablos I, Wildner S, Asam C, Wallner M, Gadermaier G (2016) Pollen allergens for molecular diagnosis. *Current Allergy and Asthma Reports* 16:31. <https://doi.org/10.1007/s11882-016-0603-z>
- Peberdy JF (1980) *Developmental Microbiology*. Boston, MA: Springer.
- Pfavayi L, Sibanda E, Mutapi F (2020) The pathogenesis of fungal-related diseases and allergies in the African population: The state of the evidence and knowledge gaps. *International Archives of Allergy and Immunology* 181:257-269. <https://doi.org/10.1159/000506009>
- Riario-Sforza G, Marinou A (2017) Hypersensitivity pneumonitis: a complex lung disease. *Clinical and Molecular Allergy* 15(1):6. <https://doi.org/10.1186/s12948-017-0062-7>
- Roper M, Pepper R, Brenner M, Pringle A (2008) Explosively launched spores of ascomycete fungi have drag-minimizing shapes. *Proceedings of the National Academy of Sciences* 105:20583-20588. <https://doi.org/10.1073/pnas.0805017105>
- Ryan M, Clark C (2015) Allergic fungal rhinosinusitis and the unified airway: the role of antifungal therapy in AFRS. *Current Allergy and Asthma Reports* 15(12):75. <https://doi.org/10.1007/s11882-015-0573-6>
- Sabit M, Wong C, Andaya A, Ramos JD (2020) Pollen allergen skin test and specific IgE reactivity among Filipinos: A community-based study. *Allergy, Asthma and Clinical Immunology* 16(1):74. <https://doi.org/10.1186/s13223-020-00471-9>
- Sánchez-Borges M, Martin BL, Muraro AM, Wood RA, Agache IO, Ansotegui IJ, Casale TB, Fleisher TA, Hellings PW, Papadopoulos NG, Peden DB, Sublett JL, Tilles SA, Rosenwasser L (2018) The importance of allergic disease in public health: an iCAALL statement. *The World Allergy Organization Journal* 11(1):8. <https://doi.org/10.1186/s40413-018-0187-2>
- Schnyder B, Pichler WJ (2009) Mechanisms of drug-induced allergy. *Mayo Clinic Proceedings* 84:268-272. [https://doi.org/10.1016/S0025-6196\(11\)61145-2](https://doi.org/10.1016/S0025-6196(11)61145-2)
- Schütze N, Lehmann I, Bönisch U, Simon J, Polte T (2010) Exposure to mycotoxins increases the allergic immune response in a murine asthma model. *American Journal of Respiratory and Critical Care Medicine* 181:1188-1199. <https://doi.org/10.1164/rccm.200909-1350OC>
- Shah A, Panjabi C (2014) Allergic aspergillosis of the respiratory tract. *European Respiratory Review*

- 23:8-29. <https://doi.org/10.1183/09059180.00007413>
- Sheldon J, Wheeler R, Riches P (2014) Immunology for clinical biochemists. In: Marshall W, Lapsley M, Day A, Ayling R (eds.) *Clinical Biochemistry: Metabolic and Clinical Aspects*, 560-603. <https://doi.org/10.1016/b978-0-7020-5140-1.00030-4>
- Simon-Nobbe B, Denk U, Pöll V, Rid R, Breitenbach M (2007) The spectrum of fungal allergy. *International Archives of Allergy and Immunology* 145:58-86. <https://doi.org/10.1159/000107578>
- Songnuan W, Bunnag C, Soontrapa K, Pacharn P, Wangthan U, Siri wattanakul U, Malainual N (2018) Airborne fungal spore distribution in Bangkok, Thailand: correlation with meteorological variables and sensitization in allergic rhinitis patients. *Aerobiologia* 34:513-524. <https://doi.org/10.1007/s10453-018-9527-5>
- Sun S, Heitman J (2011) Is sex necessary? *BMC Biology* 9:1-4. <https://doi.org/10.1186/1741-7007-9-56>
- Vijayakumar R, Aboody M, Alturaiki W, Alsagaby S, Sandle T (2017) A study of airborne fungal allergens in sandstorm dust in Al-Zulfi, Central region of Saudi Arabia. *Journal of Environmental and Occupational Science* 6:27-33. <https://doi.org/10.5455/jeos.20170120094512>
- Virella G, Hipp W, John Jr. J, Kahaleh B, Ford M, Fudenberg H (1979) Nephelometric detection of soluble immune complexes: Methodology and clinical applications. *International Archives of Allergy and Immunology* 58:402-410. <https://doi.org/10.1159/000232219>
- Walsh T, Hayden R, Larone D (2018) *Larone's Medically Important Fungi: A Guide to Identification*. USAS: ASM Press.
- Wang L, Lin X (2012) Morphogenesis in fungal pathogenicity: Shape, size, and surface. *PLOS Pathogens* 8:e1003027. <https://doi.org/10.1371/journal.ppat.1003027>
- Watanabe T (2010) *Pictorial atlas of soil and seed fungi*. Boca Raton, Florida: CRC Press.
- Whicher J, Warren C, Chambers R (1984) Immunochemical assays for immunoglobulins. *Annals of Clinical Biochemistry: International Journal of Laboratory Medicine* 21:78-91. <https://doi.org/10.1177/000456328402100202>
- Whitehead KA, Liauw CM, Lynch S, El Mohtadi M, Amin M, Preuss A, Deisenroth T, Verran J (2021) Diverse surface properties reveal that substratum roughness affects fungal spore binding. *iScience* 24:102333. <https://doi.org/10.1016/j.isci.2021.102333>
- Woodfolk JA, Commins SP, Schuyler AJ, Erwin EA, Platts-Mills TA (2015) Allergens, sources, particles, and molecules: Why do we make IgE responses? *Allergy International* 64:295-303. <https://doi.org/10.1016/j.alit.2015.06.001>
- Yamamoto N, Bibby K, Qian J, Hospodsky D, Rismani-Yazdi H, Nazaroff WW, Peccia J (2012) Particle-size distributions and seasonal diversity of allergenic and pathogenic fungi in outdoor air. *The ISME Journal* 6:1801-1811. <https://doi.org/10.1038/ismej.2012.30>
- Żukiewicz-Sobczak WA (2013) The role of fungi in allergic diseases. *Postepy Dermatologii i Alergologii* 30:42-45. <https://doi.org/10.5114/pdia.2013.33377>

